

**PATIENT INFORMATION**

(Complete the following or send patient demographic sheet.)  
Please fax copy of patient's insurance card including both sides

Name: \_\_\_\_\_  
DOB: \_\_\_\_\_ Gender: M / F Height: \_\_\_\_\_ Wt: \_\_\_\_\_  
Address: \_\_\_\_\_  
City, State, Zip: \_\_\_\_\_  
Home Phone: \_\_\_\_\_  
Alternate Phone: \_\_\_\_\_  
Last 4 digits SS#: \_\_\_\_\_ Email: \_\_\_\_\_

**PRESCRIBER INFORMATION**

Prescriber Name: \_\_\_\_\_  
State Lic. #: \_\_\_\_\_ NPI: \_\_\_\_\_  
Facility Name: \_\_\_\_\_  
Address: \_\_\_\_\_  
City, State, Zip: \_\_\_\_\_  
Phone: \_\_\_\_\_ Fax: \_\_\_\_\_  
Contact: \_\_\_\_\_  
Phone: \_\_\_\_\_ Email: \_\_\_\_\_

**CLINICAL INFORMATION – STATEMENT OF MEDICAL NECESSITY**

K50.90 Crohn's disease NOS     K51.90 Ulcerative Colitis     Other: \_\_\_\_\_  
Please indicate current or previous treatments and treatment duration below:  
 NSAIDS    Duration: \_\_\_\_\_     Corticosteroids    Duration: \_\_\_\_\_  
 Methotrexate    Duration: \_\_\_\_\_     Azathioprine    Duration: \_\_\_\_\_  
 Sulfasalazine    Duration: \_\_\_\_\_     5 – ASA    Duration: \_\_\_\_\_  
 6 – MP    Duration: \_\_\_\_\_     Other:    Duration: \_\_\_\_\_  
Other medications patient is currently taking: \_\_\_\_\_  
TB/PPD Test given?     Yes     No    Date: \_\_\_\_\_ Results: \_\_\_\_\_ Allergies: \_\_\_\_\_  
BSA: \_\_\_\_\_  
Expected date of first/next dose: \_\_\_\_\_ Date of last dose: \_\_\_\_\_

**PRESCRIPTION INFORMATION**

Medication	Directions	Quantity	Refills
Cimzia® <input type="checkbox"/> Starter Kit <input type="checkbox"/> Prefilled Syringe <input type="checkbox"/> Lyophilized Powder	<input type="checkbox"/> Initial dose: Inject 400 mg SC at week 0, week 2, and week 4 <input type="checkbox"/> Maintenance dose: Inject 400 mg SC every 4 weeks	1 starter kit	0
		28 day supply	
Entyvio® <input type="checkbox"/> 300 mg vial	<input type="checkbox"/> Initial Dose: Infuse 300mg IV over 30 minutes at week 0, week 2 and week 6 <input type="checkbox"/> Maintenance dose: Infuse 300mg IV over 30 minutes every 8 weeks	Initial dose	
		QS	
Humira® <input type="checkbox"/> Starter Kit <input type="checkbox"/> 40 mg Pen <input type="checkbox"/> 40 mg PFS <input type="checkbox"/> Starter Kit CF <input type="checkbox"/> 40 mg CF Pen <input type="checkbox"/> 40 mg CF Prefilled Syringe	<input type="checkbox"/> Initial Dose: Inject 160 mg SC on day 1, 80 mg on day 15, then 40 mg thereafter beginning on day 29 <input type="checkbox"/> Inject 40 mg SC every other week <input type="checkbox"/> Inject 40 mg SC once a week	1 starter kit	0
		28 day supply	
Remicade® <input type="checkbox"/> 100 mg vial	<input type="checkbox"/> Initial Dose: Infuse _____ mg/kg (_____ mg) IV at weeks 0, 2, and 6 weeks. <input type="checkbox"/> Maintenance Dose: Infuse _____ mg/kg (_____ mg) IV every _____ weeks.	QS	
Rinvoq® <input type="checkbox"/> Starter Kit: 45 mg <input type="checkbox"/> 15 mg <input type="checkbox"/> 30 mg	<input type="checkbox"/> Initial Dose: Take 45mg po once a day for 8 weeks. <input type="checkbox"/> Maintenance Dose: Take 15mg po once a day. <input type="checkbox"/> Maintenance Dose: Take 30mg po once a day.	28	1
		30	
Simponi® <input type="checkbox"/> 100 mg SmartJect <input type="checkbox"/> 100 mg PFS	<input type="checkbox"/> Initial Dose: Inject 200mg SC at week 0, then 100mg at week 2, then maintenance dose <input type="checkbox"/> Maintenance Dose: Inject 100mg SC once every 4 weeks	3 units	0
		28 day supply	
Stelara® <input type="checkbox"/> 130 mg vial <input type="checkbox"/> 90mg Prefilled Syringe	Initial Dose: Infuse IV over at least 1 hour. <input type="checkbox"/> ≤ 121 lbs 260 mg (2 vials) <input type="checkbox"/> > 121 lbs to 187 lbs 390 mg (3 vials) <input type="checkbox"/> > 187 lbs 520 mg (4 vials) Maintenance Dose: Inject 90 mg SC 8 weeks after initial IV infusion, then every 8 weeks thereafter	QS	0
		QS	
Skyrizi® <input type="checkbox"/> 600 mg vial <input type="checkbox"/> 360 mg PF on-body injector	<input type="checkbox"/> Initial Dose: Infuse 600 mg IV over at least 1 hour on week 0, 4 and 8. <input type="checkbox"/> Maintenance Dose: Inject 360 mg SC at week 12, and every 8 weeks thereafter.	1	2
		1	
Xeljanz® <input type="checkbox"/> 5 mg tabs <input type="checkbox"/> 10 mg tabs	<input type="checkbox"/> Initial Dose: Take 10 mg PO BID for 8 weeks <input type="checkbox"/> Maintenance Dose: Take 5 mg PO BID <input type="checkbox"/> Maintenance Dose: Take 10 mg PO BID		
Xifaxan® <input type="checkbox"/> 550mg Tablet	<input type="checkbox"/> Take 1 tablet by mouth 3 times a day	42	0
Zeposia® <input type="checkbox"/> 7 Day Titration Pack <input type="checkbox"/> 0.92 Capsules	<input type="checkbox"/> Initial Dose: Take as directed on titration pack. <input type="checkbox"/> Maintenance Dose: Take 1 capsule by mouth once a day beginning day 8.	1	
		30	
Other:			

Ship Medications To:     Physician's Clinic     Patient's Home

Injection Training Needed:     Yes     No

Prescriber Signature \_\_\_\_\_ Date \_\_\_\_\_ Brand Name Required?     Yes

I authorize this pharmacy and its representatives to act as my authorized agent to secure coverage and initiate the insurance prior authorization process for my patient and to sign any necessary forms on my behalf as my authorized agent, including the receipt of any required prior authorization forms, financial treatment and the receipt and submission of patient lab values and other patient data. In the event that this pharmacy determines that it is unable to fulfill this prescription, I further authorize this pharmacy to forward this information and any related materials related to coverage of this product to another pharmacy of the patient's choice or in the patient's insurer's provider network.